

Is the 2-Flavor Chiral Transition of First Order?

Sanatan Digal

*Department of Physics, University of Tokyo,
Tokyo 113-0033, Japan*

*Radiation Lab., RIKEN, 2-1 Hirosawa,
Wako, Saitama, 351-0198, Japan*

and

*Institute of Mathematical Sciences,
C.I.T. Campus, Taramani,
Chennai 600 113, India*

We study the effects of the interaction between the Chiral condensate and the Polyakov loop on the chiral transition within an effective Lagrangian. We find that the effects of the interaction change the order of the phase transition when the explicit breaking of the Z_N symmetry of the Polyakov loop is large. Our results suggest that the chiral transition in 2-flavor QCD may be first order.

PACS numbers: PACS numbers: 12.38.Mh

I. INTRODUCTION

QCD matter with N_f flavors, N colors and zero baryon chemical potential undergoes two finite temperature phase transitions, the chiral transition and the deconfinement. The chiral transition is associated with the spontaneous breaking of the chiral symmetry, $SU(N_f)_L \times SU(N_f)_R \rightarrow SU(N_f)_V$, below the critical temperature (T_χ) for massless quarks. The order parameter for this transition is the chiral condensate. On the other hand the deconfinement transition is associated with the spontaneous symmetry breaking, $Z_N \subset SU(N) \rightarrow \mathbf{1}$, above the critical temperature T_d for infinitely heavy quarks. The order parameter for this transition is the Polyakov loop expectation value. For finite non-zero quark masses both the chiral symmetry and the Z_N symmetry are explicitly broken. Nevertheless these two transitions show up as crossover, second or first order transitions depending on the values of the quark masses. The chiral transition depends on the number of quark flavors and the deconfinement transition depends on the number of colors. Since the chiral symmetry and the Z_N symmetry seem nothing to do with each other one would expect that these two transitions occur independently. However lattice QCD calculations have shown that both these transitions occur simultaneously, $T_\chi = T_d$ [1-5]. Furthermore strong correlation between the chiral condensate and the Polyakov loop is observed around the phase transition point [6]. This is clear evidence that there is interaction between these two order parameters. So studying the interaction between the chiral condensate and the Polyakov loop is fundamental to understanding the interplay between the chiral transition and deconfinement. There are several studies on the possible causes of the simultaneous occurrence of the chiral transition and the deconfinement transition. Mixing between the gauge and matter field operators has been suggested to explain the simultaneous chiral and deconfinement transitions [7, 8]. Some other studies consider that for small quark masses the chiral transition drives the deconfinement transition [6, 9].

Though most of the studies are concerned with why $T_\chi = T_d$ only a few have considered the effect of the interaction on the phase transition itself [10]. It seems natural to expect that if the interaction between the two order parameters can result in the simultaneous occurrence of the two transitions then the interaction may also have important effect on the phase transition. One of the most interesting cases to study for the possible effects of this interaction is the 2-flavor chiral transition. Lattice QCD calculations have not yet been able to settle on the order of this phase transition.

Lattice calculations by different groups do not agree on the order of this phase transition. Some lattice groups find the transition is second order [3] and other groups find the transition is first order [11]. Conventionally this transition is believed to be second order and in the universality class of $O(4)$ Heisenberg magnet [12]. But the effect of interaction between the chiral order parameter (Φ) and the Polyakov loop (L) may change the behavior of this transition. So in the present work we investigate the effect of interaction between Φ and L on the 2-flavor chiral transition within an effective Lagrangian.

Previously, the effect of interaction between the chiral order parameter and the Polyakov loop has been studied in the renormalization group approach [10]. The main difference between the present work and previous studies is that we consider the explicit breaking of the Z_N symmetry. Explicit breaking of the Z_N symmetry can be introduced in the effective Lagrangian by terms such as $\sim (L + L^\dagger)$, $\sim (L + L^\dagger)\Phi^\dagger\Phi$. Previous studies [10] have considered interaction term, such as $(LL^\dagger)(\Phi^\dagger\Phi)$, which respects both the chiral and the Z_N symmetry. However, in the chiral limit, the interaction terms need not respect the Z_N symmetry. So terms such as $\sim (L + L^\dagger)\Phi^\dagger\Phi$ should be considered. As we discuss later such a interaction term is always present if the explicit Z_N symmetry breaking is large, for example in the chiral limit. For simplicity we consider $N = 2$ color QCD. We expect that different N will not qualitatively change the physics we are discussing here. For $N_f = 2$ the chiral order parameter Φ is a four component vector field whereas for $N = 2$ the Polyakov loop L is a real scalar field. In this work we basically study the effect of the three terms, L , $L\Phi^\dagger\Phi$ and $L^2\Phi^\dagger\Phi$ in the effective Lagrangian. Our main result is that the strong explicit breaking of the Z_N symmetry can make the chiral transition first order. We show that the chiral transition can be first order even at the mean field level. We also carry out numerical Monte Carlo simulations which confirm the first order phase transition for large enough explicit Z_N symmetry breaking. We mention here that the $N_f = 2$ chiral transition can be first order from the interaction term $L^2\Phi^\dagger\Phi$ without Z_N symmetry breaking [13]. However lattice QCD results indicate that effect of the Z_N symmetric interaction term is small. A possible first order chiral transition can result more likely from the explicit Z_N symmetry breaking as we will argue later.

Conventionally explicit symmetry breaking weakens a phase transition. But our results suggest that for a system with two order parameter fields explicit symmetry breaking can make the transition stronger. It is interesting to note that the chiral order parameter Φ does not couple to gauge fields directly. The gauge fields seem to affect the chiral phase transition indirectly, through the Polyakov loop. We mention here that the effect of explicit symmetry breaking discussed here should not be restricted to the Z_N symmetry. We expect that the explicit breaking of chiral symmetry may have some effect on the deconfinement transition in the large quark mass region. We mention here that interaction between the chiral order parameter and the diquark fields is considered to study the chiral/color-superconducting transition at low temperature and high density [14].

This paper is organized as follows. In section-II we describe the effective Lagrangian and discuss the effect of the interaction between Φ and L on the chiral transition. We describe our numerical Monte Carlo calculations and results in section-III. The discussions and conclusions are presented in section-IV.

II. THE EFFECTIVE LAGRANGIAN AND THE EFFECT OF THE BROKEN Z_N SYMMETRY

We consider the following Lagrangian [9] in 3 dimension for the Φ and the L fields,

$$\begin{aligned}\mathcal{L} &= \frac{1}{2}|\nabla\Phi|^2 + \frac{1}{2}(\nabla L)^2 + V(\Phi, L), \\ V(\Phi, L) &= \frac{m_\Phi^2}{2}|\Phi|^2 + \frac{\lambda_\Phi}{4}|\Phi|^4 + \frac{m_L^2}{2}L^2 + \frac{\lambda_L}{4}L^4 - gL^2|\Phi|^2 - cL|\Phi|^2 - eL\end{aligned}\quad (1)$$

Some of the parameters of this reduced 3D Lagrangian depend explicitly on temperature. This Lagrangian is invariant under $O(4)$ rotation of the Φ field. The last two terms of $V(\Phi, L)$ break the Z_2 symmetry ($L \rightarrow -L$) explicitly. The interactions between the chiral order parameter and the Polyakov loop are taken care by 5th and 6th terms in $V(\Phi, L)$.

The signs of the couplings g and c decides the correlation between the fluctuations of the Φ and the L fields. For example when $c > 0$, the thermal average of the correlation between the Φ and L fluctuations, $\langle(\delta L)(\delta|\Phi|)\rangle$, is positive. When $c < 0$, $\langle(\delta L)(\delta|\Phi|)\rangle$ is negative. Such "anti-correlation" is seen between the fluctuations of the the chiral condensate and the Polyakov loop in the results of lattice QCD calculations[6]. The correlation between the variations of the two order parameters with respect to temperature is of the same sign as the correlation between the fluctuations. Note that in the above Lagrangian when $c \neq 0$ the Φ field acts like an ordering field for the L field. In the chiral symmetric phase the chiral order parameter is small and the Polyakov loop expectation value is large. The large expectation value of L , in the chiral symmetric phase, can result only from the last term in $V(\Phi, L)$ (with $e > 0$) because $|\Phi|$ is small.

As we have mentioned before previous studies [10] have considered only the interaction term $gL^2|\Phi|^2$. When the coupling parameters $c = 0 = e$, the chiral transition and the deconfinement transition do not always occur simultaneously. For large value of the coupling g the transitions can occur simultaneously and are of first order when coefficients of $|\Phi|^2$ and L^2 are negative in Eq.1 [13, 14]. A large positive g would increase the critical temperature for the deconfinement transition as the coefficient of L^2 term is negative for high temperatures. Lattice results on the other hand show that inclusion of dynamical quarks decrease the deconfinement transition temperature. So in QCD the coupling g should be small. For small g both the transitions are second order and do not occur simultaneously.

The coupling parameters c and e represent the strength of the Z_N explicit breaking. So they should increase with decreasing quark masses as the Z_N breaking becomes severe. This can be seen explicitly in the large quark mass region [15]. To see the effect of the explicit Z_N breaking let us consider $g = 0 = e$ and $c \neq 0$. At the mean field level one can consider the temperature variation of the parameters m_Φ^2 and m_L^2 . For simplicity we fix $\lambda_{\phi, L} > 0$, $m_L^2 > 0$ and vary the m_Φ^2 parameter. To find the m_Φ^2 dependence of L and Φ expectation values one has to solve the following coupled equations,

$$\begin{aligned}\lambda_\Phi|\Phi|^3 + m_\Phi^2|\Phi| - 2c|\Phi|L &= 0 \\ \lambda_LL^3 + m_L^2L - c|\Phi|^2 &= 0.\end{aligned}$$

We have checked numerically that for large enough c these equations give two solutions which correspond to a degenerate minima of the effective potential $V(\Phi, L)$ at some particular value of m_Φ^2 . It may seem surprising that the potential $V(\Phi, L)$ has degenerate minima even though there is no cubic term for Φ and L in it. However because of the coupling c these two fields are mixed. Though the mixing angle varies as m_Φ^2 is varied. With variation of m_Φ^2 the minimum of $V(\Phi, L)$, in the $|\Phi| - L$ plane, moves in directions other than the $|\Phi|$ or L axes. To understand how the minimum of $V(\Phi, L)$ behaves one should express $V(\Phi, L)$ in terms of variables which are the linear combinations of $|\Phi|$ and L . This would invariably lead to cubic terms of the new fields in the effective potential. For some choice of parameters the cubic term may then be important to cause degenerate

minima of $V(\Phi, L)$. Even if the explicit symmetry breaking is not strong enough at mean field level fluctuations at higher order can make the transition first order. At one loop the fluctuations of the Φ field will contribute to a non zero 3-point function of the L field. This 3-point function can be calculated perturbatively. In the high temperature approximation the 3-point function is given by,

$$\sim T \frac{c^3}{m^3}, \quad (2)$$

assuming zero momentum to all the external L -lines. m here is the mass of the Φ field fluctuations. Given a suitable choice of parameters in the effective Lagrangian the three point function can be significant. The consequence of this is a $\alpha(T)L^3$ term in the potential $V(\Phi, L)$ with temperature dependent $\alpha(T)$. This term can cause discontinuous change in L as temperature varies. When L field changes discontinuously the Φ field will also changes discontinuously because of the coupling term $cL\Phi^2$. If the coefficient of the Φ^2 term changes from $+ve$ to $-ve$ due to discontinuous change in L then Φ will change discontinuously from zero to non-zero. The L^3 term can also come from other types of explicit symmetry breaking coupling terms but we think $cL\Phi^2$ is the simplest term in our model.

The situation with $c = 0$ and $g \neq 0 \neq e$ is same as the one discussed above. When the explicit symmetry breaking parameter e is large, Z_2 symmetry of the L field is lost and the L field always has non-zero expectation value L_0 . In order to study the fluctuations one must expand the potential $V(\Phi, L)$ around L_0 , $L = L_0 + \bar{L}$. This gives rise to a term like $gL_0\bar{L}\Phi^2$ coming from the expansion of $gL^2\Phi^2$ around $L = L_0$. Now the term $gL_0\bar{L}\Phi^2$ is similar to the one discussed above. So there can be first order transition when $g \neq 0 \neq e$ like in the case when $c \neq 0$. We observed that at the mean field the transition becomes second order for large quartic couplings λ_Φ . Now in the following section we discuss the numerical Monte Carlo simulations of the effective Lagrangian (Eq.1) and the results. These calculations include higher order as well as non perturbative fluctuations.

III. NUMERICAL CALCULATIONS AND RESULTS

The numerical Monte Carlo calculations of the the model Eq.1 is done by discretizing the action $S = \int \mathcal{L} d^3x$ on a 3 dimensional N_s^3 lattice. We employ the following rescaling of the fields variables and the couplings,

$$\begin{aligned} \Phi &\rightarrow \frac{\Phi\sqrt{\kappa_\Phi}}{a}, \lambda_\Phi \rightarrow \frac{\lambda_\Phi}{\kappa_\Phi^2}, m_\Phi^2 \rightarrow \frac{2 - 4\lambda_\Phi - 6\kappa_\Phi}{\kappa_\Phi a^2}, \\ L &\rightarrow \frac{L\sqrt{\kappa_L}}{a}, \lambda_L \rightarrow \frac{\lambda_L}{\kappa_L^2}, m_L^2 \rightarrow \frac{2 - 4\lambda_L - 6\kappa_L}{\kappa_L a^2}, \\ g &\rightarrow \frac{g}{\kappa_\Phi \kappa_L}, c \rightarrow \frac{c}{a\sqrt{\kappa_L \kappa_\Phi}}, e \rightarrow \frac{e}{a^3\sqrt{\kappa_L}}. \end{aligned} \quad (3)$$

a is the lattice spacing and κ_Φ and κ_L are the hopping parameters for the Φ and the L fields respectively. After the rescaling of the fields and the coupling parameters the discretized lattice

action becomes,

$$\begin{aligned}
S = \sum_x & - \kappa_\Phi \sum_\mu \Phi_x \Phi_{x+\mu} + |\Phi_x|^2 + \lambda_\Phi (|\Phi_x|^2 - 1)^2 \\
& - \kappa_L \sum_\mu L_x L_{x+\mu} + L_x^2 + \lambda_L (L_x^2 - 1)^2 \\
& - g L_x^2 |\Phi_x|^2 - c L_x |\Phi_x|^2 - c L_x
\end{aligned} \tag{4}$$

Here $\Phi_x(L_x)$ represents the value of the $\Phi(L)$ field at the lattice site x . $x + \mu$ represents the six nearest neighbor lattice sites to x . We adopted the pseudo heat-bath method used for the Higgs updating in SU(2)+Higgs studies [16]. To update Φ_x at a lattice site x we write the probability distribution $P(\Phi_x)$ of Φ_x as,

$$\begin{aligned}
P(\Phi_x) & \sim \text{Exp}[-S_1(\Phi_x) - S_2(\Phi_x)], \quad \text{with} \\
S_1(\Phi_x) & = \alpha \left(\Phi_x - \frac{A}{2\alpha} \right)^2, \quad S_2(\Phi_x) = \lambda_\Phi (|\Phi_x|^2 - B)^2 \\
A & = \kappa_\Phi \sum_\mu \Phi_{x+\mu}, \quad B = 1 - \frac{1}{2\lambda_\Phi} + \frac{\alpha}{2\lambda_\Phi} + \frac{cL_x + gL_x^2}{2\lambda_\Phi}
\end{aligned} \tag{5}$$

The coefficient α is a parameter chosen so that we get a reasonable acceptance rate for the new Φ_x . Once α is chosen, a Gaussian random number is generated according to the distribution $\text{Exp}[-S_1(\Phi_x)]$. Then this random number is accepted as the new value of Φ_x with the probability $\text{Exp}[-S_2(\Phi_x)]$. Using this procedure Φ_x is updated at all the lattice sites. Then we do the updating of L_x along the same steps. The process of updating Φ_x and L_x on the entire lattice is repeated about 20 times between successive measurements. We measure the magnetizations



$$M_\Phi = \frac{1}{V} \sum_x \Phi_x, \quad M_L = \frac{1}{V} \sum_x L_x, \tag{6}$$

where $V = N_s^3$ is the number of lattice sites. The expectation values of the Φ and the L fields are given by the thermal averages (average over the measurements), $\langle \Phi \rangle = \langle |M_\Phi| \rangle$ and $\langle L \rangle = \langle M_L \rangle$. We take the absolute value of M_Φ for $\langle \Phi \rangle$ because a normal average of M_Φ is usually not a well behaved observable.



The numerical simulations were carried out on a $N_s = 16$ lattice. In this work we do not intent to explore the phase diagram of the model (Eq.1) but to show that for suitable choice of parameters the phase transition can change from second order to first order. Here we present results for two sets of parameters. For one set we fix the couplings $g = e = 0$ and for the second set we fix the coupling $c = 0$. For the first set of parameters we choose, $\lambda_\Phi = 0.004$, $\lambda_L = 0.0020$, $\kappa_L = 0.01$ and $c = 0.1$. We observe the hysteresis of $\langle \Phi \rangle$ and $\langle L \rangle$ by varying the parameter κ_Φ . In FIG.1 we show the hysteresis loop of $\langle \Phi \rangle$ and in FIG.2 we show the hysteresis loop of $\langle L \rangle$.

Since we take c to be positive we see $\langle \Phi \rangle$ and $\langle L \rangle$ increase or decrease simultaneously. The choice of the values for the parameters is such that the variation of $\langle \Phi \rangle$ and $\langle L \rangle$ are similar in magnitude. For $c < 0$ increase in $\langle \Phi \rangle$ should lead to decrease in $\langle L \rangle$. So in this case the hysteresis loop for $\langle \Phi \rangle$ will look somewhat similar to Fig.1 while the hysteresis loop for $\langle L \rangle$ will be more or less inverted about the y-axis.

For the second set of parameters we choose $\lambda_\Phi = 0.0055$, $\lambda_L = 0.0010$, $\kappa_L = 0.14$, $g = -0.02$ and $e = 0.9$. The hysteresis loop of $\langle \Phi \rangle$ and $\langle L \rangle$ are observed by again varying the parameter κ_Φ . The choice of κ_L and e is such that the expectation value of L is non-zero and positive. The sign of g

FIG. 1: The hysteresis of $\langle \Phi \rangle$ vs κ_Φ FIG. 2: The hysteresis of $\langle L \rangle$ vs κ_Φ

assures that increase in $\langle \Phi \rangle$ leads to decrease in $\langle L \rangle$ and vice-versa as the parameter κ_Φ is varied. The hysteresis curves of the two order parameters are shown in FIG.3 and FIG.4. The values of $\lambda_{\Phi,L}$ in our calculations are chosen so that we can see first order phase transition clearly and the variation of $\langle \Phi \rangle$ and $\langle L \rangle$ are of order $O(1)$ across the transition point. Note that with suitable choice of κ_L and λ_L one change the average of the Polyakov loop across the transition point. The choice g was such that the chiral transition turned out to be second order when the coupling e was set to zero. We also did simulations on a 4D lattice. The results in this case are very similar to the 3D simulations.

FIG. 3: The hysteresis of $\langle \Phi \rangle$ vs κ_Φ FIG. 4: The hysteresis of $\langle L \rangle$ vs κ_Φ

The results in FIG.1-4 show strong first order phase transition for the Φ and L fields. By fixing the coupling parameters g , c and e we observed that the strength of the transition depends on the values of the quartic couplings λ_Φ and λ_L . The transition becomes weaker with increase in any of the quartic couplings $\lambda_{\Phi,L}$. However even for larger quartic couplings a suitable choice of the parameters g , c and e made the transition strong first order.

We also did calculations with small explicit symmetry breaking for the Φ field by considering a linear Φ term in the Lagrangian. In this case we found that the hysteresis loops of both $\langle\Phi\rangle$ and $\langle L\rangle$ shrinking in size with increase in the coefficient of linear Φ term in the Lagrangian. These results suggest that the transition becomes weaker when the chiral symmetry is explicitly broken.

IV. DISCUSSION AND CONCLUSIONS

Using a simple effective Lagrangian, which captures the chiral symmetry and Z_N symmetry of QCD, we have investigated the effect of the explicit symmetry breaking on the chiral phase transition. Since we consider the chiral limit the Z_2 symmetry of the Polyakov loop is explicitly broken. As we incorporate the explicit Z_2 breaking interaction terms in the effective Lagrangian we find the chiral transition becomes first order. We observed that the first order transition becomes weaker when a small explicit symmetry breaking is considered for the chiral order parameter Φ . The results of our calculations show that two flavor chiral transition is of first order for large enough Z_N explicit breaking. As we have mentioned before at present some lattice studies suggest that the transition is first order [11] and some other studies show the transition is second order [3]. These conflicting results may be because the quark masses studied are not small enough or the lattices used are not big enough.

In the previous study, for $N_f = 2$ and $N = 3$ [10], the second order chiral transition becomes first order when coupled to the Polyakov loop. This is because the deconfinement transition is first order for $N = 3$ with a cubic term $\sim (L^3 + L^{*3})$ in the effective potential with exact Z_3 symmetry. However when the quark masses are finite and decrease the deconfinement transition becomes weaker. This can be understood due to explicit breaking of Z_3 symmetry. Already in the large quark mass region the deconfinement transition becomes crossover which implies the explicit symmetry breaking dominates over the effects of the above Z_3 symmetric cubic term. For smaller quark masses the explicit Z_3 symmetry breaking likely grows and expected to be maximal in the chiral limit. So for smaller quark masses, i.e in the chiral limit one should rather consider the effect of the explicit breaking of the Z_3 symmetry. The effects of explicit symmetry breaking discussed in this work should not be restricted to the explicit breaking of the Z_N symmetry. For 2-flavor and 2-color QCD the deconfinement transition in the large quark mass region may have the effects coming from the explicit breaking of the chiral symmetry. It may be possible that this effect change the phase transition behavior of the deconfinement transition in the heavy quark mass region.

Acknowledgments

We have benefited greatly from discussions with T. Hatsuda and A. M. Srivastava. We would like to thank G. Baym, S. Datta, M. Laine, T. Matsuura and M. Ohtani for comments. This work is supported by the JSPS Postdoctoral Fellowship for Foreign Researchers.

-
- [1] J. B. Kogut *et al.*, Phys. Rev. Lett.**50**, 393 (1983).
 - [2] M. Fukugita and A. Ukawa, Phys. Rev. Lett.**57**, 503 (1986).
 - [3] F. Karsch, and E. Laermann, Phys. Rev. **D50**, 6954 (1994).
 - [4] S. Aoki *et al.*, Phys. Rev.**D57**, 3910 (1998).
 - [5] F. Karsch, E. Laermann, and A. Peikert, Nucl. Phys. **B605**, 579 (2001).
 - [6] S. Dital, E. Laermann, and H. Satz, Eur. Phys. J. **C18**, 583, (2001).
 - [7] F. Karsch, E. Laermann, and C. Schmidt, Phys. Lett. **B520**, 41 (2001).
 - [8] Y. Hatta and K. Fukushima, Phys. Rev. **D69**, 097502 (2004).

- [9] A. Mocsy, F. Sannino, and K. Tuominen, Phys. Rev. Lett. **92**, 182302, (2004); *ibid*, JHEP **0403**, 044 (2004).
- [10] S. Midorikawa, H. So, and S. Yoshimoto, Z. Phys. **C34**, 307, (1987).
- [11] M. Fukugita, H. Mino, M. Okawa, and A. Ukawa, Phys. Rev. **D42**, 2936 (1990); M. D'Elia, A. D. Giacomo, and C. Pica, Phys. Rev. **D72**, 114510 (2005).
- [12] K. Rajagopal and F. Wilczek, Nucl. Phys.**B399**, 395, (1993); R. Pisarski and F. Wilczek, Phys. Rev. **D29**, 338 (1984).
- [13] P. M. Chaikin and T. C. Lubensky, Principles of Condensed Matter Physics, Combridge Univ. press, (Combridge, 1995).
- [14] G. Baym, T. Hatsuda, M. Tachibana and N. Yamamoto, private communication.
- [15] F. Green and F. Karsch, Nucl. Phys. **B238**, 297, (1984).
- [16] B. Bunk, Nucl. Phys. (Proc. Suppl.) **42**, 566, (1995).